



**Susquehanna River
Basin Commission**

**Susquehanna Large River
Assessment Project**

Publication 265
September 2009

Report by Tyler Shenk
Water Quality Specialist

www.srbc.net

SRBC • 1721 N. Front Street • Harrisburg, PA 17102 • 717.238.0423 • srbc@srbc.net

Printed on recycled paper

ABSTRACT

In 2002, the Susquehanna River Basin Commission (SRBC) conducted a pilot study to determine appropriate methods of biologically assessing the large rivers of the Susquehanna River Basin (basin). Based on the results of that survey, SRBC determined that a combination of rock basket samplers and traditional Rapid Bioassessment Protocol (RBP) methods was the most efficient and consistent collection method to sample the Susquehanna River. These methods were implemented in the 2005 Susquehanna Large River Assessment Project (Hoffman, 2006) at 25 stations on the mainstem Susquehanna River and at the mouths of its major tributaries: the West Branch Susquehanna River, the Juniata River, and the Chemung River.

The U.S. Environmental Protection Agency (USEPA) has developed a field operations manual for the National River and Stream Assessment (NRSA), detailing data collection methods for both wadeable and nonwadeable streams (USEPA, 2008). In 2007, SRBC adapted this protocol for the 25 stations previously sampled (Hoffman, 2008). In late summer 2008, SRBC staff collected data at 17 of the same stations following the NRSA protocols.

Composite benthic macroinvertebrate samples were collected at each station from three D-frame net sweeps at each of ten transects. Field and laboratory water quality samples and overall observations of the site also were collected at each site.

Two of the sites were designated as nonimpaired, ten sites were slightly impaired, and five sites were moderately impaired. Only 6.9 percent (28 of 408) of the water quality values exceeded their respective limits, indicating fairly good water quality in the Susquehanna River.

SRBC plans to continue to use the adapted NRSA river protocols in the future. Fish collection may be added to the protocol in subsequent years, and SRBC plans to develop protocols to properly assess the reservoirs near the mouth of the Susquehanna River.

INTRODUCTION

SRBC has been performing biological assessments throughout the basin since the late 1970s. When USEPA introduced the first version of the RBP manual (Plafkin and others, 1989), SRBC adopted those methods for use in its interstate stream monitoring program and its rotating subbasin surveys. However, neither the previous nor current RBP methods (Barbour and others, 1999) used by SRBC in the aforementioned surveys accurately depict the biological integrity of the basin's large rivers: the mainstem Susquehanna, Chemung, West Branch Susquehanna, and Juniata Rivers. Thus, in 2002, SRBC initiated a pilot project to determine proper methods of biologically assessing the large rivers in the basin. From this pilot project, staff determined that a combination of rock-filled basket samplers and traditional RBP methods was the most effective and consistent collection method for sampling the Susquehanna River (Hoffman, 2003).

In summer 2005, SRBC staff collected biological and water quality data at 25 stations on the mainstem Susquehanna River and at the mouth of its major tributaries using the methodology described above. In 2007, staff changed the methodology to mimic the methods drafted by USEPA for the NRSA (USEPA, 2008). These methods have been used for the past two years.

Although the NRSA data collection includes fish, physical habitat, toxicology, and other parameters in addition to

benthic macroinvertebrates, SRBC staff chose to focus efforts on benthic macroinvertebrate sampling. Benthic macroinvertebrates were used to assess biological conditions for several reasons. Benthic macroinvertebrates are sensitive to a wide range of stressors, have a wide range of documented pollution tolerances, and are found in a wide variety of habitats throughout lotic systems (Flotemersch and others, 2001a). Additionally, SRBC has background macroinvertebrate data from various sites on the large rivers of the basin from subbasin surveys and interstate streams monitoring, as well as the previous river assessment studies.

Geography

The Susquehanna River Basin is the largest river basin on the east coast of the United States, draining 27,510 square miles. The Susquehanna River originates at Otsego Lake, N.Y., and flows 444 miles through New York, Pennsylvania, and Maryland to the Chesapeake Bay at Havre de Grace, Md.

This year's Large River Assessment stretched from Great Bend to Marietta, Pa., and encompassed a total of 17 stations: two in the Upper Susquehanna Subbasin; nine in the Middle Susquehanna Subbasin; one at the mouth of the West Branch Susquehanna River; four in the Lower Susquehanna Subbasin; and one at the mouth of the Juniata River. Downstream of Marietta, Pa., the river flows through a series of dams and reservoirs, which this protocol is not designed to assess.



Collecting macroinvertebrate samples at Wyalusing, Pa.